

Math 60670 Homework 4

Due Monday, February 24.

Problem 1: Let ∇ be a torsion-free linear connection, and ω a 1-form. Show that

$$d\omega(X, Y) = (\nabla_X\omega)(Y) - (\nabla_Y\omega)(X)$$

for any $X, Y \in \mathcal{X}(M)$.

Problem 2: Let ∇ be a linear connection. Given a curve $\gamma(t) : I \rightarrow M$ and a vector $V \in T_{\gamma(s)}M$ (for some $s \in I$), let us write $P_{\gamma, V}(t)$ for the parallel transport of V along γ with respect to ∇ . If $X, Y \in \mathcal{X}(M)$ and $p \in M$, show that

$$\nabla_X Y|_p = \lim_{t \rightarrow 0} \frac{1}{t} (P_{\gamma, Y(\gamma(t))}(0) - Y(0))$$

for any curve $\gamma : (-\epsilon, \epsilon) \rightarrow M$ satisfying $\gamma(0) = p$, $\gamma'(0) = X$.

Problem 3: Let (M, g) be an oriented Riemannian manifold. Recall that if ω is a k -form on M , and X is a vector field, then $\iota_X\omega$ is the $(k-1)$ -form obtained by contracting against X in the first slot:

$$(\iota_X\omega)(V_1, \dots, V_{k-1}) := \omega(X, V_1, \dots, V_{k-1}).$$

The divergence operator $\text{div} : \mathcal{X}(M) \rightarrow C^\infty(M)$ is defined by

$$d(\iota_X dV) = \text{div}(X)dV.$$

A) Show that if M is a compact manifold-with-boundary, then

$$\int_M \text{div}(X)dV = \int_{\partial M} \langle X, N \rangle dV_{\partial M},$$

where $dV_{\partial M}$ is the volume-form on ∂M with the inducted metric and orientation, and N is the outwards pointing conormal of ∂M .

B) Show that if $\phi \in C^\infty(M)$, then div satisfies the following product rule:

$$\text{div}(\phi X) = \langle \text{grad}\phi, X \rangle + \phi \text{div}(X).$$

C) Using the above or otherwise, show that in local coordinates div can be written as

$$\text{div}(X) = \frac{1}{\sqrt{g}} \partial_i (\sqrt{g} X^i), \quad g = \det(g_{ij}).$$

D) Show that if ∇ is the Levi-Civita connection, then

$$\text{div}(X) = \text{tr}(\nabla X) \equiv (\nabla X)_i^i \equiv \sum_i \langle e_i, \nabla_{e_i} X \rangle,$$

(e_i being any an ON basis of the tangent space).